

Title: The effects of repetitive transcranial magnetic stimulation on empathy: A systematic review and meta-analysis

Yang, C. C., Khalifa, N. & Vollm, B. A. (2017). The effects of repetitive transcranial magnetic stimulation on empathy: A systematic review and meta-analysis. *Psychological Medicine*, 48 (5), pp.737-750.

Link to repository:

<https://repository.nottinghamshirehealthcare.nhs.uk/handle/123456789/2239>

Additional information:

This article has been published in a revised form in *Psychological Medicine* <http://doi.org/10.1017/S003329171700232X>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © Cambridge University Press.

Publisher: Cambridge University Press

Version note:

The version presented here may differ from the published version or from the version of record. If you wish to cite the following item, it is advised to consult the publisher version. Access to the publisher version can be found via the repository URL listed above.

For more information about this article, or the research repository, please contact repository@nottshc.nhs.uk

Please cite the published version

Nottinghamshire Healthcare NHS Foundation Trust
Institutional Repository
repository.nottinghamshirehealthcare.nhs.uk

**The effects of repetitive transcranial magnetic stimulation on empathy: A
systematic review and meta-analysis**

Cheng-Chang Yang, Najat Khalifa, Birgit Völlm*

**Division of Psychiatry and Applied Psychology, School of Medicine, University
of Nottingham**

Corresponding author

*Address for correspondence: Professor Birgit Völlm, Division of Psychiatry and
Applied Psychology, School of Medicine, University of Nottingham, UK
(Email: birgit.vollm@nottingham.ac.uk)

Financial support: N/A

Word count: 4878

Abstract

Empathy is a multi-dimensional concept with affective and cognitive components, the latter often referred to as Theory of Mind (ToM). Impaired empathy is prevalent in people with neuropsychiatric disorders, such as personality disorder, psychopathy, and schizophrenia, highlighting the need to develop therapeutic interventions to address this. Repetitive transcranial magnetic stimulation (rTMS), a non-invasive therapeutic technique that has been effective in treating various neuropsychiatric conditions, can be potentially used to modulate empathy. To our knowledge, no systematic reviews or meta-analyses in this field have been conducted. The aim of the current study was to review the literature on the use of rTMS to modulate empathy in adults. Seven electronic databases (AMED, Cochrane library, Embase, Medline, Pubmed, PsycInfo, and Web of Science) were searched using appropriate search terms. Twenty-two studies were identified, all bar one study involved interventions in healthy rather than clinical populations, and 18 of them, providing results for 24 trials, were included in the meta-analyses. Results showed an overall small, but statistically significant, effect in favour of active rTMS in healthy individuals. Differential effects across cognitive and affective ToM were evident. Subgroup analyses for cognitive ToM revealed significant effect sizes on excitatory rTMS, offline paradigms, and non-randomised design trials. Subgroup analyses for affective ToM revealed significant effect sizes on excitatory rTMS, offline paradigms, and non-randomised design trials. Meta-regression revealed no significant sources of heterogeneity. In conclusion, rTMS may have discernible effects on different components of empathy. Further research is required to examine the effects of rTMS on empathy in clinical and non-clinical populations, using appropriate empathy tasks and rTMS protocols.

The effects of repetitive transcranial magnetic stimulation on empathy: A systematic review and meta-analysis

Successful human socialisation is heavily influenced by the abilities to detect and understand cognitive and emotional processes in others. These abilities are referred to as the Theory of Mind (ToM) and empathy (Gallese, 2003; Young *et al.*, 2010; Keuken *et al.*, 2011; Krall *et al.*, 2016). Clinicians and researchers use these terms interchangeably, but there is no universal consensus on their definitions and constructs. For example, some authors regard empathy as a two-component construct with affective and cognitive components (e.g., Reniers *et al.*, 2011) whilst others (e.g., Blair, 2005) have proposed a three-component construct by adding a motor component to reflect the act of mirroring the motor responses of the observed person (motor empathy). Some commentators view cognitive empathy as synonym to ToM which is the ability to attribute mental states, such as desires, intentions and beliefs, to others (Frith & Frith, 1999). Some authors have favoured a ToM model with two distinct components, namely affective and cognitive (e.g., Kalbe *et al.*, 2010). Others have suggested that empathy and ToM encompass similar underlying abilities that are discernible at the neural level (e.g., Reniers *et al.*, 2014). More recently, Dvash & Shamay-Tsoory (2014) argued in favour of a two-component construct of empathy, namely emotional and cognitive empathy (also referred to as ToM), with distinct neuroanatomical underpinnings (Fig.1). According to this model cognitive empathy (ToM) has two distinct subcomponents, namely affective ToM and cognitive ToM.

Several brain regions have been implicated in cognitive ToM, including medial prefrontal cortex (mPFC), dorsolateral prefrontal cortex (DLPFC), temporoparietal

junction (TPJ) and temporal poles (Frith & Frith, 1999; Völlm *et al.*, 2006; Carrington & Bailey, 2009; Reniers *et al.*, 2014). Brain areas implicated in the regulation of affective ToM include mPFC, particularly the ventral portion (Shamay-Tsoory & Aharon-Peretz, 2007; Shamay-Tsoory *et al.*, 2009; Sebastian *et al.*, 2012), inferior frontal gyrus (IFG), anterior cingulate cortex, and amygdala (Shamay-Tsoory *et al.*, 2009; Gonzalez-Liencre *et al.*, 2013; Gentili *et al.*, 2015).

Self-report inventories commonly used to measure empathy include the Hogan Empathy Scale (Hogan, 1969), the Interpersonal Reactivity Index (IRI; Egger *et al.*, 1997), the Balanced Emotional Empathy Scale (BEES; Mehrabian, 2000), the Empathy Quotient (EQ; Behan *et al.*, 2015), and the Questionnaire of Cognitive and Affective Empathy (QCAE; Reniers *et al.*, 2011). Behavioural measures of cognitive empathy (ToM) are primarily performance-based and include such tasks as first-order (Baron-Cohen *et al.*, 1985) and second-order false-belief (Baron-Cohen, 1989) tasks for assessing cognitive ToM, the Reading the Mind in the Eyes (RMET) for evaluating affective ToM (Baron-Cohen *et al.*, 2001), and the Faux Pas Recognition (FPR) test (Stone *et al.*, 1998) and the Yoni task (Shamay-Tsoory & Aharon-Peretz, 2007) for assessing both affective and cognitive ToM.

Impairment of social functioning consequent upon impaired empathy has been reported in a range of neuropsychiatric conditions, including psychopathy, antisocial personality disorder (Dolan & Fullam, 2004), schizophrenia (Bragado-Jimenez & Taylor, 2012), major depressive disorder (MDD; Schreiter *et al.*, 2013), autistic spectrum disorder (ASD; Shimoni *et al.*, 2012), temporal lobe epilepsy (Li *et al.*, 2013), Alzheimer's disease (Laisney *et al.*, 2013), Parkinson's disease (Yu *et al.*,

2012), and other neurodegenerative diseases (Poletti *et al.*, 2012). Empathy is highly correlated with violence (Jolliffe & Farrington, 2004) and plays a pivotal role in the violence inhibition system (Blair *et al.*, 2005). Thus, enhancement of empathy has been regarded as a major treatment goal in criminogenic programmes (Day *et al.*, 2010; Reidy *et al.*, 2013). However, conventional psychological interventions for empathy enhancement have proved less effective in certain offender groups, particularly those with psychopathy (Reidy *et al.*, 2013), highlighting the need to develop alternative therapeutic interventions to enhance empathy, of which transcranial magnetic stimulation (TMS), especially its repetitive format (rTMS), is an example (Glenn & Raine, 2008; Glannon, 2014).

TMS is a non-invasive technique used to deliver brief, high-intensity magnetic pulses to the brain inducing localised neuronal depolarization to regulate cortical excitability that underlies the modulation of cortical networks (Luber & Lisanby, 2014). In general, high frequency (≥ 5 Hz) rTMS and its newer version, intermittent theta burst stimulation (iTBS), facilitate cortical excitability, whereas low frequency (about 1 Hz) rTMS and continuous theta burst stimulation (cTBS) contribute to opposite effects (Pascual-Leone *et al.*, 2000; Huang *et al.*, 2005; Wassermann & Zimmermann, 2012). rTMS has been used to treat a variety of neurological and psychiatric diseases (see Wassermann & Zimmermann, 2012) and to enhance cognitive functions in healthy volunteers (see Hsu *et al.*, 2015) and in people with MDD (Serafini *et al.*, 2015). Table S1 provides more information about the effects of TMS in clinical populations. Additionally, rTMS has been used to modulate empathy with some promising effects (see Hetu *et al.*, 2012; Schuwerk *et al.*, 2014a). However, findings are inconsistent likely due to differences in the tasks used to measure

empathy, experimental designs, targeted brain regions, and rTMS parameters, including the paradigms used (i.e., online or offline), stimulus intensity (measured as a percentage of resting motor threshold [rMT] or of maximum stimulator output [MSO]), frequency and number of pulses.

We therefore aimed to conduct a systematic review and meta-analysis of the literature on the effects of rTMS on empathy in healthy and clinical populations to integrate the evidence base and to determine if certain TMS parameters or brain regions selected are associated with stronger effects on specific domains of empathy. Whilst effective interventions involving healthy individuals could potentially be extended to clinical populations, as we shall describe later in this review, all the studies included in this review, bar one study, involved interventions in healthy groups. Due to the overlaps between the concepts of empathy and ToM, in this review we have conceptualised empathy in accordance with the model proposed by Dvash & Shamay-Tsoory (2014) as outlined above. We followed PRISMA-P guidelines (Moher *et al.*, 2015; Shamseer *et al.*, 2015) in the reporting of this review where applicable.

Method

Data sources

Using the terms "transcranial magnetic stimulation" or "TMS" combined with "theory of mind", "ToM", "empathy", "mentalizing", "role taking", or "perspective taking", a systematic search of the literature on the effects of TMS on empathy was conducted on 25 May 2016 of seven electronic databases (AMED, Cochrane library, Embase, Medline, PsycInfo, Pubmed, Web of Science). The International Clinical Trials Registry Platform (World Health Organization), Dissertation Abstracts, Google, and the library catalogues of the University of Nottingham were also searched to identify grey literature in the field. No filters were added regarding the age of study participants, publication time or language of publication (see online supplement Table S2 for search syntax). References of eligible articles were searched manually for potentially eligible studies missed by the electronic searches.

Study selection

Empirical studies were included in the review if they: (1) involved adult participants without dementia or other major neurological conditions; (2) used rTMS as an active intervention; (3) had a comparison group or control condition; and (4) used behavioural tasks to assess empathy. Of the 508 papers originally identified, 22 met the inclusion criteria (see Fig. S1 and Table S3) and were quality assessed using the quality assessment tool for quantitative studies (National Collaborating Centre for Methods and Tools, 2008) on the domains of selection bias, study design, confounders, blinding, data collection method, withdrawals and dropouts, intervention integrity, and statistical analyses.

Of the 22 studies included in the review, four (Uddin *et al.*, 2006; Balconi *et al.*, 2010; Hoekert *et al.*, 2010; Lev-Ran *et al.*, 2012) were excluded from the meta-analyses due to lack of sufficient data to allow effect size calculation and only after exhausting attempts to obtain this information from the authors.

Data extraction and analyses

A standardised form was used to extract information concerning authors, study objectives, sample characteristics, inclusion/exclusion criteria, study design, experimental processes, rTMS protocols, outcome variables, and analytic strategy.

We originally intended to conduct separate meta-analyses of studies involving clinical populations and healthy individuals using the random-effects model and, where applicable, in accordance with the model proposed by Dvash & Shamay-Tsoory (2014) with its components: cognitive empathy (i.e., ToM, including cognitive ToM and affective ToM) and affective empathy. However, this has not been possible due to there being only one study in the field (Enticott *et al.*, 2014). Therefore, the meta-analyses presented in this review include only studies involving healthy subjects. Measures of cognitive ToM included the cognitive component of the Yoni task, moral judgement, false-belief tasks, and action-understanding tools. Measures of affective ToM included the RMET, tasks of facial expression recognition, the affective component of the Yoni task, affective go/no-go tasks, the faux pas test and emotional egocentricity. While it can be argued that facial expression recognition is not a test of empathic abilities, the model proposed by Dvash & Shamay-Tsoory (2014) regards emotional recognition as a component of affective ToM. This view has been supported by other commentators (e.g., Poletti *et al.* 2012), Therefore,

tasks measuring emotional recognition, such as facial expression recognition tasks, were included in the review.

Effect size was regarded as positive if the active rTMS effect was in the predicted direction and negative if it was in the opposite direction. Moreover, when a study entailed multiple stimulation sites, each trial of the different stimulation sites was used as the unit of analysis for the purpose of meta-analysis. A pooled effect size was used if a study provided multiple outcomes (e.g., accuracy and reaction time, score of each subscale, or short-term and long-term performance). Only the comparison between experimental and sham group (condition) was selected when a trial consisted of more than one control group or condition (e.g., one group receiving rTMS at a control site and another receiving sham stimulation). Effect sizes represented as Hedges' g and 95% confident intervals (CI) were calculated according to the differences between experimental (real stimulation) and control (sham stimulation) conditions in post-stimulation evaluations or "online" performance divided by pooled standard deviation.

The Q and I^2 statistics (Higgins & Thompson, 2002; Higgins *et al.*, 2003) were used to assess consistency between studies. The Q statistic represents the level of heterogeneity while the I^2 index specifies the total variation from between-study variance. A P value $\leq .05$ and an I^2 value of greater than 40% were deemed as indicative of moderate heterogeneity. Funnel plots (Egger & Smith, 1995), the Egger test (Egger *et al.*, 1997), and Begg and Mazumdar rank correlation tests (Begg & Mazumdar, 1994) were used to test for the presence of a potential publication bias.

In cases where publication bias was evident, the Trim and Fill procedure (Egger & Smith, 1995) was applied to correct it.

In order to identify variables which could contribute to alternation of empathy, pre-specified subgroup analyses were performed with the unit of trial by merging the data according to the rTMS parameters, including effect (“excitatory” vs. “inhibitory”), stimulation paradigm (“online” vs. “offline”), study design (“randomised” vs. “non-randomised”), stimulation site and task of outcome measurement.

Meta-regression was employed to examine the impact of between-study variation on study effect sizes. The effect size from each trial was set as the dependent variable while age, gender, intensity of stimulation, total pulses per condition, and weighted number of pulses (i.e., total number of rTMS pulses multiplied by intensity) were selected as predictor variables. All the quantitative analyses were performed using Stata 13.1 (StataCorp, 2013).

Results

Study characteristics

Table 1 summaries study characteristics. In summary, 22 studies involving 466 participants (82% males; mean age: 24.45 years; range: 18-59 years) were included in the review. For studies recruiting participants from clinical populations, there was only one study (Enticott *et al.*, 2014), recruiting patients with ASD as subjects.

Sixteen of the included studies were conducted in Europe, three in North America (Uddin *et al.*, 2006; Young *et al.*, 2010; Keuken *et al.*, 2011), two in Australia (Krause *et al.*, 2012; Enticott *et al.*, 2014) and one in Israel (Lev-Ran *et al.*, 2012). The most common study design employed was non-randomised crossover ($n = 15$), allocating the sequence of intervention conditions with counterbalancing ($n = 10$) or unspecified ($n = 5$) method. Of the six studies randomly allocating participants, two (Keuken *et al.*, 2011; Enticott *et al.*, 2014) were parallel randomised controlled trials and the other four (Costa *et al.*, 2008; Kalbe *et al.*, 2010; Giardina *et al.*, 2011; Lev-Ran *et al.*, 2012) were randomised crossover trials. The remaining one between-subject study (Silani *et al.*, 2013) did not mention the method of participant allocation.

Various tasks were used to assess empathy, including facial expression recognition tasks with materials derived from Ekman & Friesen (1976), the RMET or its modified version, the Yoni task, scenarios using video clips assessing individuals' capability of social judgement or action-understanding, the false belief task and the faux pas task. With regard to published self-report instruments, only one study (Enticott *et al.*, 2014) selected a self-report measure, the IRI, as the empathy measure. The number of pulses within each experimental session ranged from 120 to 3000. The majority of the reviewed studies ($n = 15$) set the intensity of the pulses to 100% or more of rMT,

while other four studies used subthreshold intensity (Costa *et al.*, 2008; Hoekert *et al.*, 2010; Giardina *et al.*, 2011; Michael *et al.*, 2014). The remaining three studies (Young *et al.*, 2010; Keuken *et al.*, 2011; Krall *et al.*, 2016) selected MSO as the index of intensity. The DLPFC, mPFC (ventral or dorsal portion), TPJ, and IFG were targeted as the main sites for stimulation. The most common control condition was vertex stimulation ($n = 11$). Five studies did not report the detail of their sham protocol.

Quality assessment

Of the twenty-two studies included, only one study (Enticott *et al.*, 2014) attracted a rating of “strong”, nineteen studies were rated as “moderate”, and two studies as “weak” (Table S4). Poor rating on selection bias was the most common reason for not reaching the “strong” quality threshold. The two weak ratings were due to vulnerability to confounders (Silani *et al.*, 2013) and poor description of the reliability and validity of the outcome measures used (Michael *et al.*, 2014). For rTMS reproducibility, most of the reviewed studies ($n = 16$) provided all necessary parameters, but two studies (Balconi *et al.*, 2010; Silani *et al.*, 2013) failed to provide information in relation to the type of coil utilised and four studies (Pobric & Hamilton, 2006; Costa *et al.*, 2008; Balconi *et al.*, 2011; Balconi & Bortolotti, 2012) lacked comprehensive information about the duration of the intervention. Only three studies described adverse effects relating to the administration of rTMS, with one study indicating no adverse effects observed (Young *et al.*, 2010) and the other two studies reporting minor post-rTMS side effects (Enticott *et al.*, 2014) and one syncope event (Kalbe *et al.*, 2010).

Meta-analysis

Effects of rTMS on empathy in clinical populations

Since there was only one trial (Enticott *et al.*, 2014) involving participants with a mental disorder it was not possible to conduct a meta-analysis to examine the rTMS effect on empathy in clinical populations. This study (Enticott *et al.*, 2014) showed that deep high frequency rTMS applied bilaterally to the dorsal mPFC in patients with ASD did not have a statistically significant facilitatory effects on empathy ($g = -0.22$; 95% CI, -1.55 to -0.01, $p = 0.016$), cognitive empathy ($g = -0.32$; 95% CI, -1.07 to 0.44, $p = 0.41$), or affective empathy ($g = 0.08$; 95% CI, -0.66 to 0.82, $p = 0.21$).

Effects of rTMS on empathy in healthy volunteers

Twenty-four trials extracted from reports of 17 studies were included for the meta-analysis of the effects of rTMS on empathy. This revealed a significant small overall effect size ($g = 0.29$; 95% CI, 0.10 to 0.48, $p = 0.003$) as plotted in Fig. 2a. A moderate level of heterogeneity was observed across the studies ($Q_{23} = 39.22$, $p = .019$; $I^2 = 41.4\%$). Separate meta-analyses were conducted for trials involving cognitive empathy with its two components; cognitive and affective ToM. However, it was not possible to conduct a meta-analysis on the effects of rTMS on affective empathy due to lack of studies in the field.

Effects of rTMS on cognitive ToM

The meta-analysis of findings from 16 trials on the effects of rTMS on cognitive ToM showed a non-significant mean effect ($g = 0.12$, 95% CI, -0.15 to 0.40, $p = .39$; see also Fig. 2b). The trim and fill procedure applied suggested an estimated mean effect size of -0.13 after imputing five missing trials (Fig. S2b). A moderate heterogeneity

was found across trials ($Q_{16} = 30.64$, $p = .01$; $I^2 = 51.0\%$). The funnel plot was asymmetrical by visual inspection (Fig. S2a), but neither the Begg's test ($z = 0.95$, $p = .34$) nor the Egger's test (intercept₁₆ = 2.42, $t = 1.18$, 2-tailed $p = .26$) suggested publication bias.

The subgroup analyses (Table 2) revealed a non-significant mean effect for inhibitory rTMS ($g = 0.03$, 95% CI, -0.27 to 0.33, $p = .83$) but a significant one for excitatory rTMS ($g = 0.58$, 95% CI, 0.05 to 1.10, $p = .03$). For the stimulation paradigm, since all trials with offline paradigms applied inhibitory rTMS and all trials with online paradigms applied excitatory rTMS, the results of the subgroup analysis were the same (offline: $g = 0.03$, 95% CI, -0.27 to 0.33, $p = .83$; online: $g = 0.58$, 95% CI, 0.05 to 1.10, $p = .03$). Moreover, the subgroup analysis for study designs revealed a non-significant mean effect size for trials with randomised design ($g = -0.16$, 95% CI, -0.56 to 0.25, $p = .45$) but a significant one for trials with non-randomised design ($g = 0.40$, 95% CI, 0.13 to 0.67, $p = .004$). Furthermore, the subgroup analysis for stimulation sites revealed non-significant mean effect sizes for all stimulation sites, including TPJ ($g = 0.26$, 95% CI, -0.04 to 0.56, $p = .09$), DLPFC (including IFG) ($g = -0.09$, 95% CI, -0.71 to 0.53, $p = .79$) and mPFC ($g = 0.04$, 95% CI, -0.44 to 0.52, $p = .87$). Finally, the subgroup analysis for the nature of outcome measure tasks revealed non-significant mean effect sizes for false-belief tasks ($g = 0.10$, 95% CI, -0.21 to 0.41, $p = .51$) and intention attribution tasks ($g = -0.10$, 95% CI, -0.57 to 0.37, $p = .69$) but a significant large mean effect size for action-understanding tasks ($g = 0.82$, 95% CI, 0.34 to 1.30, $p = .001$).

The meta-regression analysis across trials showed that none of between-study variables significantly predicted the effects of rTMS (mean age of participants: $\beta = 0.08$, $p = .55$; gender ratio: $\beta = -1.01$, $p = .11$; intensity of stimulation: $\beta = -0.03$, $p = .26$; number of pulses per condition: $\beta = -0.005$, $p = .45$; weighted number of pulses: $\beta = 0.005$, $p = .48$).

Effects of rTMS on affective ToM

The meta-analysis of results from 15 trials on the effects of rTMS on affective ToM showed a significant small mean effect ($g = 0.26$, 95% CI, 0.02 to 0.50, $p = .03$) with a moderate heterogeneity ($Q_{14} = 25.98$, $p = .03$; $I^2 = 46.1\%$; see also fig. 2c). The funnel plot (Fig. S3a) and the Egger's test (intercept₁₇ = -4.39, $t = -2.55$, 2-tailed $p = .02$) showed evidence of publication bias. However, the Begg's test ($z = 1.48$, $p = .14$) and the trim and fill procedure did not show evidence of publication bias.

Further subgroup analyses showed that the mean effect size of inhibitory rTMS trials failed to reach statistical significance ($g = 0.25$, 95% CI, -0.00 to 0.51, $p = .052$). It was not possible to calculate the mean effect size for excitatory rTMS since there was only one trial (Balconi & Canavesio, 2013) in this subgroup which showed a positive effect ($g = 0.33$). For stimulation paradigms, trials with "offline" paradigms revealed a non-significant mean effect ($g = 0.10$, 95% CI, -0.12 to 0.32, $p = .35$) while trials with "online" paradigm showed a significant moderate effect ($g = 0.52$, 95% CI, 0.05 to 1.00, $p = .03$). The subgroup analysis for study design revealed a non-significant mean effect size for trials with randomised design ($g = -0.06$, 95% CI, -0.36 to 0.24, $p = .71$) but a significant one for trials with non-randomised design ($g = 0.43$, 95% CI, 0.123 to 0.73, $p = .006$). Regarding the sites of stimulation, all three

locations revealed non-significant mean effect sizes (TPJ: $g = -0.14$, 95% CI, -0.74 to 0.46, $p = .65$; DLPFC [including IFG]: $g = 0.28$, 95% CI, -0.35 to 0.91, $p = .39$; mPFC: $g = 0.22$, 95% CI, -0.07 to 0.52, $p = .14$). For type of measurement, the mean effect sizes for trials using emotion recognition tasks ($g = 0.32$, 95% CI, -0.06 to 0.69, $p = .10$) and faux-pas recognition tasks ($g = -0.08$, 95% CI, -0.50 to 0.35, $p = .73$) were not significant.

The meta-regression analysis across trials showed that none of between-study variables significantly predicted the effects of rTMS (mean age of participants: $\beta = 0.07$, $p = .44$; gender ratio: $\beta = -0.68$, $p = .22$; intensity of stimulation: $\beta = 0.15$, $p = .07$; number of pulses per condition: $\beta = 0.02$, $p = .11$; weighted number of pulses: $\beta = -0.02$, $p = .11$).

Discussion

This study aimed to examine the literature on the effects of rTMS on empathy and, where relevant, to determine which intervention parameters were associated with stronger effects. Our findings show that rTMS has a significant but small overall effect on empathy in healthy participants and that this effect varied according to empathy domains, cognitive or affective ToM. It has not been possible to draw valid conclusions regarding the effect of rTMS on empathy in clinical population as there was only one study conducted in the field.

The meta-analysis of rTMS studies relating to cognitive ToM revealed a non-significant effect size indicating that rTMS may not be effective in modulating cognitive ToM. Moreover, the results suggested that there might be five unpublished trials investigating this issue with negative findings. In contrast, a significant effect size was found on the meta-analysis of rTMS studies for affective ToM though the magnitude of effect was small. These findings of dissimilar effects of rTMS support the idea of examining subcomponents of empathy separately as they are associated with distinct brain regions (Dvash & Shamay-Tsoory, 2014).

Our subgroup analyses further identified parameters associated with a positive effect of rTMS, including excitatory vs. inhibitory rTMS and online vs. offline paradigms. However, these finding should be interpreted with caution due to the relatively small number of trials, particularly for excitatory rTMS. Although previous studies (e.g., Robertson *et al.*, 2003) suggest that the duration of the rTMS after-effect only persists for half of the stimulation time, physiological evidence indicates that the rTMS aftereffect decays gradually with time (Eisenegger *et al.*, 2008). Nevertheless,

given that completion of conventional tasks measuring empathy is time-consuming, it is less likely to detect significant rTMS effect on empathy from experiments with offline paradigm.

Surprisingly, the subgroup analysis by stimulation site did not reveal statistically significant mean effects across different brain regions pertaining to specific empathetic components. The literature suggests differential roles of specific brain regions: The dorsal part of mPFC and TPJ (particularly the right side) for cognitive ToM (e.g., Denny *et al.*, 2012) and the ventral part of mPFC and IFG for affective ToM (Sebastian *et al.*, 2012; Dal Monte *et al.*, 2014). It would thus be expected to find significant effects if rTMS is administered to these regions, but not to other regions. However, we found no significant effect applying rTMS to TPJ for cognitive ToM or IFG for affective ToM and only one included trial (Keuken *et al.*, 2011) explored affective ToM targeting at these crucial regions (e.g., IFG), a firm conclusion cannot be drawn at this stage. It is worth noting here that *the issue of spatial resolution is an inherent limitation of TMS research. The issue may be further compromised when non-imaging guided techniques are utilised to localise the stimulation sites.* With this in mind, and since a considerable number of studies included in this review (e.g., Balconi *et al.*, 2010; Balconi & Bortolotti, 2012; Krause *et al.*, 2012; Schuwerk *et al.*, 2014) did not utilise imaging guided techniques, we *have categorised the studies according to the effects of TMS on relatively large regions of the brain rather than smaller ones while performing subgroup analyses.* Nevertheless, the results need to be interpreted with caution.

Meta-regression revealed no differential effects in relation to participant characteristics (age, gender) or stimulation parameters (intensity, number of pulses, weighted number of pulses). This may be due to the low heterogeneity detected in relation to participants' age and gender ratio. Contrary to the findings of other meta-analytic studies (e.g., Chou *et al.*, 2015), rTMS parameters did not contribute significantly to effect sizes. A number of explanations exists as to why these findings were not replicated in this review. First, the number of studies included in this review was slightly higher than 10, the minimum number required to attain sufficient statistical power (Borenstein *et al.*, 2009). Second, the impact of the rTMS parameters may only be evident when rTMS is applied to the brain region corresponding to the task measured. Third, empathy is a multifaceted construct involving a network of brain regions, and since the effects of TMS are dose-dependent, a larger number of sessions and pulses per session may be required to modulate empathy.

Future research should examine a number of pertinent issues. For example, some of the included studies (Balconi & Bortolotti, 2013; Balconi & Canavesio, 2016) suggested that baseline level of empathy can moderate the inhibitory effect of low frequency rTMS on facial emotional recognition. Interestingly, they found people with higher levels of empathy performed better under control conditions than those with lower levels of empathy when the activity of the dorsal mPFC was inhibited. However, for the effect of facilitatory rTMS for enhancing empathetic ability, the role of baseline empathy level has not yet been investigated which is obviously a crucial issue for rTMS in clinical application. In addition, as speculated in a number of included studies, the behavioural tasks selected might not be appropriate for

outcome measures due to their low sensitivity to detect rTMS-induced effects (e.g., Keuken *et al.*, 2011; Krause *et al.*, 2012; Lev-Ran *et al.*, 2012; Enticott *et al.*, 2014; Schuwerk *et al.*, 2014b). Finally, it might be too simplistic to expect that increased excitability contributes to behavioural improvement and decreased excitability to a deterioration as others have also suggested (Sandrini *et al.*, 2011).

Strengths and limitations

A major strength of this study is that some of the studies included were relatively well designed with low dropouts rates and high reproducibility of rTMS protocols.

However, the study suffered a number of limitations in relation to selection bias, reflected by restricted participants' age range, recruitment resources and reporting adverse of effects which is essential in TMS studies (Rossi *et al.*, 2009). Further, the subgroup analysis of study design showed that more significant effects were found in non-randomised than randomised trials. This raises the question whether the results of the current study may be vulnerable to some methodological limitations. However, since a majority of included studies were rated as equivalently moderate in quality assessment, the source of heterogeneity is less likely from allocation bias and needs further investigation. While the research on rTMS application into alteration of empathy is still in its infancy, this systematic review with meta-analysis applied a broad range of search terms to enrol eligible studies with variant outcome measures and different rTMS protocols. We included both randomised and non-randomised trials as a considerable number of studies in this field used non-randomised design. Multiple databases were thoroughly searched to minimise potential publication bias. However, a number of studies could not be included in the meta-analysis due to not reporting effect sizes, outcome measures not matching our inclusion criteria and the

presence of possible publication bias. The majority of included studies applied empathy tasks providing multiple outcomes, such as accuracy and reaction time. We dealt with these multiple outcomes by averaging the effect sizes though this may have underestimated the size of effect. The number of studies included in the meta-analysis is relatively small, and this in conjunction with considerable levels of heterogeneity across the studies may have affected the power of the study. Finally, only one study involving interventions in a clinical population was included in the review and no meta-analytic data could therefore be provided for clinical samples. This highlights the urgent need to conduct clinical trials in the field.

Conclusion

The present review with meta-analysis demonstrated that rTMS has a discernible contribution to the alteration in different components of empathy although the effect sizes may not be as favourable as expected. The most encouraging finding for clinical implications is the effect of excitatory rTMS on enhancing *affective ToM*. Therefore, this review may help researchers having an interest in exploring rTMS impacts on empathy tailor their rTMS protocols to maximise its effect. Future studies in the field can potentially examine the effects of excitatory rTMS in clinical populations with impaired empathetic capabilities, such as those with ASD, psychopathy and schizophrenia. However, we do not currently know whether the same effects will be observed in these populations. rTMS parameters may have to be refined further to maximise the effects on crucial brain regions and there is a need to develop ecologically validated and sensitive empathy tasks for rTMS experiments.

Acknowledgements

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of interest

None.

References

Balconi M, Bortolotti A (2012). Detection of the facial expression of emotion and self-report measures in empathic situations are influenced by sensorimotor circuit inhibition by low-frequency rTMS. *Brain Stimulation* **5**, 330-336.

Balconi M, Bortolotti A (2013). Emotional face recognition, empathic trait (BEES), and cortical contribution in response to positive and negative cues. the effect of rTMS on dorsal medial prefrontal cortex. *Cognitive Neurodynamics* **7**, 13-21.

Balconi M, Bortolotti A, Gonzaga L (2011). Emotional face recognition, EMG response, and medial prefrontal activity in empathic behaviour. *Neuroscience Research* **71**, 251-259.

Balconi M, Canavesio Y (2013). High-frequency rTMS improves facial mimicry and detection responses in an empathic emotional task. *Neuroscience* **236**, 12-20.

Balconi M, Canavesio Y (2016). Empathy, approach attitude, and rTMS on left DLPFC affect emotional face recognition and facial feedback (EMG). *Journal of Psychophysiology* **30**, 17-28.

Balconi M, Crivelli D, Bortolotti A (2010). Detection of facial expression of emotion and self-report measures in empathic situations are influenced by ACC inhibition: rTMS evidences. *Neuropsychological Trends* **8**, 95-99.

Baron-Cohen S (1989). The autistic child's theory of mind: a case of specific developmental delay. *Journal of Child Psychology and Psychiatry and Allied Disciplines* **30**, 285-297.

Baron-Cohen S, Leslie A M, Frith U (1985). Does the Autistic-Child Have a Theory of Mind. *Cognition* **21**, 37-46.

Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I (2001). The "Reading the Mind in the Eyes" test revised version: A study with normal adults, and adults with

Asperger syndrome or high-functioning autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines* **42**, 241-251.

Begg CB, Mazumdar M (1994). Operating characteristics of a rank correlation test for publication bias. *Biometrics* **50**, 1088-1101.

Behan B, Stone A, Garavan H (2015). Right Prefrontal and Ventral Striatum Interactions Underlying Impulsive Choice and Impulsive Responding. *Human Brain Mapping* **36**, 187-198.

Blair J, Mitchell DR, Blair K (2005). *The psychopath: emotion and the brain*. Blackwell Pub.: Malden, MA.

Blair RJ (2005). Responding to the emotions of others: dissociating forms of empathy through the study of typical and psychiatric populations. *Consciousness and Cognition* **14**, 698-718.

Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009). Meta-Regression. In *Introduction to Meta-Analysis*. John Wiley & Sons: Chichester, U.K.

Bragado-Jimenez MD, Taylor PJ (2012). Empathy, schizophrenia and violence: A systematic review. *Schizophrenia Research* **141**, 83-90.

Carrington SJ, Bailey AJ (2009). Are There Theory of Mind Regions in the Brain? A Review of the Neuroimaging Literature. *Human Brain Mapping* **30**, 2313-2335.

Chou YH, Hickey PT, Sundman M, Song AW, Chen NK (2015). Effects of repetitive transcranial magnetic stimulation on motor symptoms in Parkinson disease: a systematic review and meta-analysis. *JAMA Neurology* **72**, 432-440.

Costa A, Torriero S, Oliveri M, Caltagirone C (2008). Prefrontal and temporo-parietal involvement in taking others' perspective: TMS evidence. *Behavioural Neurology* **19**, 71-74.

Dal Monte O, Schintu S, Pardini M, Berti A, Wassermann EM, Grafman J,

Krueger F (2014). The left inferior frontal gyrus is crucial for reading the mind in the eyes: Brain lesion evidence. *Cortex* **58**, 9-17.

Day A, Casey S, Gerace A (2010). Interventions to improve empathy awareness in sexual and violent offenders: Conceptual, empirical, and clinical issues. *Aggression and Violent Behavior* **15**, 201-208.

Denny BT, Kober H, Wager TD, Ochsner KN (2012). A Meta-analysis of Functional Neuroimaging Studies of Self- and Other Judgments Reveals a Spatial Gradient for Mentalizing in Medial Prefrontal Cortex. *Journal of Cognitive Neuroscience* **24**, 1742-1752.

Dolan M, Fullam R (2004). Theory of mind and mentalizing ability in antisocial personality disorders with and without psychopathy. *Psychological Medicine* **34**, 1093-1102.

Dvash J, Shamay-Tsoory SG (2014). Theory of Mind and Empathy as Multidimensional Constructs Neurological Foundations. *Topics in Language Disorders* **34**, 282-295.

Egger M, Smith GD (1995). Misleading meta-analysis. *British Medical Journal* **310**, 752-754.

Egger M, Smith GD, Schneider M, Minder C (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal* **315**, 629-634.

Eisenecker C, Treyer V, Fehr E, Knoch D (2008). Time-course of "off-line" prefrontal rTMS effects - a PET study *Neuroimage* **42**, 379-384.

Ekman P, Friesen W (1976). *Pictures of facial affect*. Consulting Psychologists Press: Palo Alto, CA.

Enticott PG, Fitzgibbon BM, Kennedy HA, Arnold SL, Elliot D, Peachey A,

Zangen A, Fitzgerald PB (2014). A double-blind, randomized trial of deep Repetitive

Transcranial Magnetic Stimulation (rTMS) for autism spectrum disorder. *Brain Stimulation* **7**, 206-211.

Frith CD, Frith U (1999). Interacting minds--a biological basis. *Science* **286**, 1692-1695.

Gallese V (2003). The roots of empathy: The shared manifold hypothesis and the neural basis of intersubjectivity. *Psychopathology* **36**, 171-180.

Gentili C, Cristea IA, Ricciardi E, Costescu C, David D, Pietrini P (2015). Neurobiological Correlates of the Attitude Toward Human Empathy. *Rivista Internazionale Di Filosofia E Psicologia* **6**, 70-87.

Giardina A, Caltagirone C, Oliveri M (2011). Temporo-parietal junction is involved in attribution of hostile intentionality in social interactions: An rTMS study. *Neuroscience Letters* **495**, 150-154.

Glannon W (2014). Intervening in the psychopath's brain. *Theoretical Medicine and Bioethics* **35**, 43-57.

Glenn A L, Raine A (2008). The neurobiology of psychopathy. *Psychiatric Clinics of North America* **31**, 463-475.

Gonzalez-Liencre C, Shamay-Tsoory SG, Brune M (2013). Towards a neuroscience of empathy: Ontogeny, phylogeny, brain mechanisms, context and psychopathology. *Neuroscience and Biobehavioral Reviews* **37**, 1537-1548.

Hetu S, Taschereau-Dumouchel V, Jackson PL (2012). Stimulating the brain to study social interactions and empathy. *Brain Stimulation* **5**, 95-102.

Higgins JP, Thompson SG (2002). Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* **21**, 1539-1558.

Higgins JP, Thompson SG, Deeks JJ, Altman DG (2003). Measuring inconsistency in meta-analyses. *BMJ* **327**, 557-560.

Hoekert M, Vingerhoets G, Aleman A (2010). Results of a pilot study on the involvement of bilateral inferior frontal gyri in emotional prosody perception: an rTMS study. *BMC Neuroscience* **11**, 93.

Hogan R (1969). Development of an Empathy Scale. *Journal of Consulting and Clinical Psychology* **33**, 307-316.

Hsu WY, Ku Y, Zanto TP, Gazzaley A (2015). Effects of noninvasive brain stimulation on cognitive function in healthy aging and Alzheimer's disease: a systematic review and meta-analysis. *Neurobiol Aging* **36**, 2348-2359.

Huang YZ, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC (2005). Theta burst stimulation of the human motor cortex. *Neuron* **45**, 201-206.

Jolliffe D, Farrington DP (2004). Empathy and offending: A systematic review and meta-analysis. *Aggression and Violent Behavior* **9**, 441-476.

Kalbe E, Schlegel M, Sack AT, Nowak DA, Dafotakis M, Bangard C, Brand M, Shamay-Tsoory S, Onur OA, Kessler J (2010). Dissociating cognitive from affective theory of mind: A TMS study. *Cortex* **46**, 769-780.

Keuken MC, Hardie A, Dorn BT, Dev S, Paulus MP, Jonas KJ, Van den Wildenberg WPM, Pineda JA (2011). The role of the left inferior frontal gyrus in social perception: An rTMS study. *Brain Research* **1383**, 196-205.

Krall SC, Volz LJ, Oberwelland E, Grefkes C, Fink GR, Konrad K (2016). The right temporoparietal junction in attention and social interaction: A transcranial magnetic stimulation study. *Human Brain Mapping* **37**, 796-807.

Krause L, Enticott PG, Zangen A, Fitzgerald PB (2012). The role of medial prefrontal cortex in theory of mind: A deep rTMS study. *Behavioural Brain Research* **228**, 87-90.

Laisney M, Bon L, Guiziou C, Daluzeau N, Eustache F, Desgranges B (2013).

Cognitive and affective Theory of Mind in mild to moderate Alzheimer's disease.

Journal of Neuropsychology **7**, 107-120.

Lev-Ran S, Shamay-Tsoory S, Zangen A, Levkovitz Y (2012). Transcranial magnetic stimulation of the ventromedial prefrontal cortex impairs theory of mind learning. *European Psychiatry* **27**, 285-289.

Li YH, Chiu MJ, Yeh ZT, Liou HH, Cheng TW, Hua MS (2013). Theory of Mind in Patients with Temporal Lobe Epilepsy. *Journal of the International Neuropsychological Society* **19**, 594-600.

Luber B, Lisanby SH (2014). Enhancement of human cognitive performance using transcranial magnetic stimulation (TMS). *Neuroimage* **85**, 961-970.

Mehrabian A (2000). Beyond IQ: Broad-based measurement of individual success potential or "emotional intelligence". *Genetic Social and General Psychology Monographs* **126**, 133-239.

Michael J, Sandberg K, Skewes J, Wolf T, Blicher J, Overgaard M, Frith CD (2014). Continuous theta-burst stimulation demonstrates a causal role of premotor homunculus in action understanding. *Psychological Science* **25**, 963-972.

Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA, PRISMA-P Group (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* **4**, 1.

National Collaborating Centre for Methods and Tools (2008). Quality Assessment Tool for Quantitative Studies Method. McMaster University: Hamilton, ON.

Pascual-Leone A, Walsh V, Rothwell J (2000). Transcranial magnetic stimulation in cognitive neuroscience--virtual lesion, chronometry, and functional connectivity. *Current Opinion in Neurobiology* **10**, 232-237.

Pobric G, Hamilton AFdC (2006). Action Understanding Requires the Left Inferior Frontal Cortex. *Current Biology* **16**, 524-529.

Poletti M, Enrici I, Adenzato M (2012). Cognitive and affective Theory of Mind in neurodegenerative diseases: Neuropsychological, neuroanatomical and neurochemical levels. *Neuroscience and Biobehavioral Reviews* **36**, 2147-2164.

Reidy DE, Kearns MC, DeGue S (2013). Reducing psychopathic violence: A review of the treatment literature. *Aggression and Violent Behavior* **18**, 527-538.

Reniers RLEP, Corcoran R, Drake R, Shryane NM, Völlm BA (2011). The QCAE: A Questionnaire of Cognitive and Affective Empathy. *Journal of Personality Assessment* **93**, 84-95.

Reniers RLEP, Völlm BA, Elliott R, Corcoran R (2014). Empathy, ToM, and self-other differentiation: An fMRI study of internal states. *Social Neuroscience* **9**, 50-62.

Robertson EM, Theoret H, Pascual-Leone A (2003). Studies in cognition: The problems solved and created by transcranial magnetic stimulation. *Journal of Cognitive Neuroscience* **15**, 948-960.

Rossi S, Hallett M, Rossini PM, Pascual-Leone A, Safety of TMS Consensus Group (2009). Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clinical Neurophysiology* **120**, 2008-2039.

Sandrini M, Umiltà C, Rusconi E (2011). The use of transcranial magnetic stimulation in cognitive neuroscience: A new synthesis of methodological issues. *Neuroscience and Biobehavioral Reviews* **35**, 516-536.

Schreier S, Pijnenborg GHM, Aan Het Rot M (2013). Empathy in adults with clinical or subclinical depressive symptoms. *Journal of Affective Disorders* **150**, 1-16.

Schuwerk T, Langguth B, Sommer M (2014a). Modulating functional and

dysfunctional mentalizing by transcranial magnetic stimulation. *Frontiers in Psychology* 5. <http://dx.doi.org/10.3389/fpsyg.2014.01309>

Schuwert T, Schecklmann M, Langguth B, Dohnel K, Sodian B, Sommer M (2014b). Inhibiting the posterior medial prefrontal cortex by rTMS decreases the discrepancy between self and other in Theory of Mind reasoning. *Behavioural Brain Research* 274, 312-318.

Sebastian CL, Fontaine NMG, Bird G, Blakemore SJ, De Brito SA, McCrory EJ P, Viding E (2012). Neural processing associated with cognitive and affective Theory of Mind in adolescents and adults. *Social Cognitive and Affective Neuroscience* 7, 53-63.

Serafini G, Pompili M, Belvederi Murri M, Respino M, Ghio L, Girardi P, Fitzgerald PB, Amore M (2015). The effects of repetitive transcranial magnetic stimulation on cognitive performance in treatment-resistant depression. A systematic review. *Neuropsychobiology* 71, 125-139.

Shamay-Tsoory SG, Aharon-Peretz J (2007). Dissociable prefrontal networks for cognitive and affective theory of mind: A lesion study. *Neuropsychologia* 45, 3054-3067.

Shamay-Tsoory SG, Aharon-Peretz J, Perry D (2009). Two systems for empathy: a double dissociation between emotional and cognitive empathy in inferior frontal gyrus versus ventromedial prefrontal lesions. *Brain* 132, 617-627.

Shamseer L, Moher D, Clarke M, Gherzi D, Liberati A, Petticrew M, Shekelle P, Stewart L A, PRISMA-P Group. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *The BMJ* 349, g7647.

Shimoni H N, Weizman A, Yoran R H, Raviv A (2012). Theory of mind, severity of

autistic symptoms and parental correlates in children and adolescents with Asperger syndrome. *Psychiatry Research* **197**, 85-89.

Silani G, Lamm C, Ruff CC, Singer T (2013). Right supramarginal gyrus is crucial to overcome emotional egocentricity bias in social judgments. *Journal of Neuroscience* **33**, 15466-15476.

StataCorp (2013). Stata Statistical Software: Release 13. College Station, TX: StataCorp LP.

Stone VE, Baron-Cohen S, Knight RT (1998). Frontal lobe contributions to theory of mind. *Journal of Cognitive Neuroscience* **10**, 640-656.

Uddin LQ, Molnar-Szakacs I, Zaidel E, Iacoboni M (2006). rTMS to the right inferior parietal lobule disrupts self-other discrimination. *Social Cognitive and Affective Neuroscience* **1**, 65-71.

Völlm BA, Taylor ANW, Richardson P, Corcoran R, Stirling J, McKie S, Deakin J FW, Elliott R (2006). Neuronal correlates of theory of mind and empathy: A functional magnetic resonance imaging study in a nonverbal task. *Neuroimage* **29**, 90-98.

Wassermann EM, Zimmermann T (2012). Transcranial magnetic brain stimulation: therapeutic promises and scientific gaps. *Pharmacology & Therapeutics* **133**, 98-107.

Young L, Camprodon JA, Hauser M, Pascual-Leone A, Saxe R (2010). Disruption of the right temporoparietal junction with transcranial magnetic stimulation reduces the role of beliefs in moral judgments. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 6753-6758.

Yu RL, Wu RM, Chiu MJ, Tai CH, Lin CH, Hua MS (2012). Advanced Theory of Mind in patients at early stage of Parkinson's disease. *Parkinsonism & Related*

Disorders **18**, 21-24.


Figures Captions



Figure 1. Empathy system adapted from Dvash and Shamay-Tsoory (2014)

Abbreviations: ACC, anterior cingulate cortex; DLPFC, dorsolateral prefrontal cortex; IFG, inferior frontal gyrus; mPFC, medial prefrontal cortex; STS, superior temporal sulcus; ToM, Theory of Mind; TPJ, temporoparietal junction; vmPFC, ventromedial prefrontal cortex

Figure 2. (a) Statistical summary and forest plot of effect sizes for empathy. **(b)** Statistical summary and forest plot of effect sizes for cognitive ToM. **(c)** Statistical summary and forest plot of effect sizes for affective ToM

Abbreviations: DLPFC, dorsolateral prefrontal cortex; ES, effect size; IFG, inferior frontal gyrus; mPFC, medial prefrontal cortex; PMC, primary motor cortex; S1, primary somatosensory area; SMG, Supramarginal gyrus; TPJ, temporoparietal junction TBS, theta burst stimulation; TPJ, temporoparietal junction








Table 1. Characteristics of included rTMS studies on empathy

Study (country)	Study design	participants number [‡] , Age(Mean± SD, range), male%, Diagnosis if not healthy volunteers	Tasks	Stimulation position	rTMS protocol (frequency, intensity, stimulation, paradigm, number of pulses per condition)	Sham method
Balconi & Bortolotti, 2012 (Italy)	UCR	18, (23.40± 2.60, 20-30), 44%	Facial expression recognition	mPFC	1Hz, 120% rMT, online, 400 pulses	Vertex stimulation & unknown sham method at FCz
Balconi & Bortolotti, 2013 (Italy)	CCR	19, (23.13± 2.11, 20-30), 47%	Facial expression recognition	dorsal mPFC	1Hz, 120% rMT, online, 400 pulses	Vertex stimulation & unknown sham method at mPFC
Balconi, Bortolotti, & Gonzaga, 2011 (Italy)	UCR	20, (23.73± 2.08, 20-30), 45%	Facial expression recognition	mPFC	1Hz, 120% rMT, online, 200 pulses	Vertex stimulation & unknown sham method at mPFC
Balconi & Canavesio, 2013 (Italy)	UCR	16, (23.11± 1.93, 20-28), 38%	Facial expression recognition	mPFC	10Hz, 120% rMT, online, 2500 pulses	Vertex stimulation & tilt (45 degree) coil at mPFC
Balconi & Canavesio, 2016 (Italy)	CCR	46, (26.77± 0.17, NA), 57%	Facial expression recognition	left DLPFC	1Hz, 120% rMT, online, 400 pulses	Vertex stimulation & tilt (45 degree) coil at left DLPFC
Balconi, Crivelli, & Bortolotti, 2010 ^c (Italy)	UCR	18, (23.46± 2.65, NA), NA	Facial expression recognition	ACC	1Hz, 120% rMT, online, 400 pulses	Vertex stimulation & unknown sham method at FCz
Bolognini et al., 2013 (Italy)	CCR	Exp1: 18, (22.6± 3.5, NA), 11% Exp2: 18, (24.5±3.8, NA), 17%	Affective go/no-go task	Exp1: right S1 Exp2: left S1	1Hz, 110% rMT, offline, 600 pulses	Exp1:left DLPFC stimulation & no stimulation Exp2: right DLPFC stimulation & no stimulation
Costa et al., 2008 (Italy)	RCR [†]	11, (22.5± 3.0, NA), 45%	Short stories: false belief/faux pas/control	left TPJ right TPJ left DLPFC right DLPFC	1Hz, 90% rMT, offline, 900 pulses	unknown sham method
Enticott et al.,	RCT	28(active: 15, sham: 13),	IRI	bilateral dorsal	5 Hz, 100% rMT, offline, 900	Sham coil

2014 (Australia)		(32.32±11.80, 18-59), 82%, ASD	RMET, Frith-Happé- animations	mPFC	pulses	
Giardina et al., 2011 (Italy)	RCR†	14, (22±3, NA), 21%	Social interaction scenarios requiring either hostile or non- hostile intentionality attributions	left TPJ right TPJ	1Hz, 90% rMT, offline, 600 pulses	Occipital cortex stimulation
Hoekert et al., 2010 ^c (Netherlands)	CCR	9, (21.8± 2.6, 18-26), 40% ^a	Emotional language task	left IFG, right IFG	5Hz, 90% rMT, online, 576 pulses	right IFG stimulation Sham coil
Kalbe et al., 2010 (Germany)	RCR†	28, (24.0± 2.7, NA), 100%	RMET, Yoni task	right DLPFC	1Hz, 100% rMT, offline, 900 pulses	Vertex stimulation
Keuken et al., 2011 (USA)	RCT†	37 (active: 18, control: 19), (20.4± 2.0, 18-29), 100%	Modified RMET, Attribution of belief and intentions; reasoning about physical causations (modified from Brunet et al., 2000)	left IFG	1Hz, 45% MSO, offline, 300 pulses	Vertex stimulation
Krall et al., 2016 (Germany)	CCR	24, (27.7± 4.5, 18 – 40), 54%	False belief task	right TPJ	cTBS, 30% MSO, offline, 600 pulses	Vertex stimulation
Krause et al., 2012 (Australia)	UCR	16, (26.42± 3.82, 18 – 40), 38%	Yoni task RMET	bilateral dorsal mPFC	1 Hz, 100% rMT, offline, 900 pulses	Sham coil
Lev-Ran et al., 2012 ^c (Israel)	RCR†	13, (24.73± 2.89, NA), 62%	Yoni task	ventral mPFC	1Hz, 100% rMT, offline, 400 pulses	Superior temporal region stimulation
Michael et al., 2014 (Denmark)	CCR	20, (23.5, 18–40), 60%	Action-understanding task	The hand and lip area in the left M1	cTBS, 70% rMT, offline, 300 pulses	Either stimulation site as control
Pobric and Hamilton, 2006 (UK)	CCR	exp1:9, (NA, 21-35), 64% ^b exp2:9, (NA, 21-35), 64% ^b	Action- understanding task	left IFG	5Hz, 110% rMT, online, 240 pulses	left occipital cortex stimulation, Vertex stimulation, &

Schuwerk et al., 2014 (Germany)	CCR	17, (22.2± 2.3, NA), 35%	False belief task requiring the computation of another's and one's own belief	posterior mPFC	1Hz, 100% rMT, offline, 2000 pulses	no stimulation Tilt (90 degree) coil at posterior mPFC
Silani et al., 2013 (Switzerland)	CCT	45 (active: 22 control: 23), (NA, NA), 0%	Judgments of pleasantness of self- or other-experienced visuo-tactile stimulation	right SMG	1Hz, 110% rMT, offline, 900 pulses	Vertex stimulation
Uddin et al., 2006 ^c (USA)	CCR	8, (26.6, NA), 25%	self–other facial discrimination task	right IPL	1Hz, 100% rMT, offline, 1200	Left IPL stimulation
Young et al., 2010 (USA)	CCR	Exp1: 8, (NA, 18-30), 38% Exp2: 12, (NA, 18-30), 42%	Moral scenarios manipulating protagonists' beliefs and action outcomes	right TPJ	Exp1: 1Hz, 70% MSO, offline, 1500 pulses Exp.2: 10Hz, 60% MSO, online, 120 pulses	5 cm posterior to the right TPJ stimulation

ACC, anterior cingulate cortex; ASD, autistic spectrum disorder; CCR, counterbalanced crossover design; CCT, clinical controlled trial; cTBS, continuous theta burst stimulation; DLPFC, dorsolateral prefrontal cortex; Exp: experiment; IFG, inferior frontal gyrus; IPL, inferior parietal lobule; IRI, Interpersonal Reactivity Index; M1, primary motor cortex; mPFC, medial prefrontal cortex; MSO, maximum of stimulator output; NA, not available; RCR, randomised crossover design; RCT, randomised controlled trial; RMET, Reading the Mind in the Eye Test; rMT, resting motor threshold; rTMS, repetitive transcranial magnetic stimulation; S1, primary somatosensory area; SMG, Supramarginal gyrus; TPJ, temporoparietal junction; UCR, crossover design with unknown allocation

† no randomisation method reported

‡ presented as number of participants included in final analysis and the number of participants in subgroups in the parenthesis

^a presented as the original sex ratio

^b presented as the sex ratio of participants in the whole study

^c not included for meta-analysis

Table 2. Subgroup analyses

	Pooled effect size			Between-study heterogeneity		
	<i>k</i>	Effect size (Hedges' <i>g</i>)	95% CI	Q test	<i>I</i> ²	<i>p</i> value
Cognitive ToM						
Total	16	0.12	-0.15-0.40	30.64	51.0%	0.010
Effect of stimulation						
Inhibitory	13	0.03	-0.27-0.33	25.66	53.2%	0.012
Excitatory	3	0.58*	0.05-1.10	1.23	0.0%	0.539
Stimulation paradigm						
Online	3	0.58*	0.05-1.10	1.23	0.0%	0.539
Off-line	13	0.03	-0.27-0.33	25.66	53.2%	0.012
Study design						
Randomised	8	-0.16	-0.56-0.25	15.83	55.8%	0.027
Non-randomised	8	0.40*	0.13-0.67	5.40	0.0%	0.611
Stimulation site						
TPJ	7	0.26	-0.04-0.56	2.50	0.0%	0.869
DLPFC (including IFG)	6	-0.09	-0.71-0.53	18.34	72.7%	0.003
mPFC	2	0.04	-0.44-0.52	0.00	0.0%	0.992
Type of used task						
False-belief	6	0.10	-0.21-0.41	1.81	0.0%	0.875
Intention attribution	7	-0.10	-0.57-0.37	16.87	64.4%	0.010
Action understanding	3	0.82*	0.34-1.30	0.18	0.0%	0.912
Affective ToM						
Total	15	0.26*	0.02-0.50	25.98	46.1%	0.026
Effect of stimulation						
Inhibitory	14	0.25	-0.00-0.51	25.97	49.9%	0.017
Excitatory	1	0.33	-0.36-1.03	-	-	-
Stimulation paradigm						
Online	5	0.52*	0.05-1.00	11.95	66.5%	0.018
Off-line	10	0.10	-0.12-0.32	6.08	0.0%	0.732
Study design						
Randomised	6	-0.06	-0.35-0.50	0.91	0.0%	0.970
Non-randomised	9	0.43*	0.12-0.73	16.71	52.1%	0.033
Stimulation site						
TPJ	2	-0.14	-0.74-0.46	0.26	0.0%	0.611
DLPFC (including IFG)	5	0.28	-0.35-0.91	19.03	79.0%	0.001
mPFC	5	0.22	-0.07-0.52	2.11	0.0%	0.716
Type of used task						
emotion recognition	8	0.32	-0.06-0.69	20.66	66.1%	0.004
faux-pas recognition	4	-0.08	-0.50-0.35	0.35	0.0%	0.950

CI, confidence interval; DLPFC, dorsolateral prefrontal cortex; IFG, inferior frontal gyrus; mPFC, medial prefrontal cortex; ToM, Theory of Mind; TPJ, temporoparietal junction

* *p* < .05

Online Supplementary materials

Table S1: rTMS effects in clinical populations (after Wassermann and Zimmermann, 2012)

Population	Effects
Depression	rTMS at DLPFC yields a medium to large effect size on reducing the severity of depressive symptoms.
Schizophrenia	Low-frequency rTMS significantly reduces intensity of auditory hallucinations but is less efficient in improving negative symptoms.
Obsessive compulsive disorder (OCD)	High-frequency rTMS may reduce compulsions; the finding has not been replicated consistently across studies.
Posttraumatic stress disorder (PTSD)	High-frequency rTMS may have positive and sustainable therapeutic effects on anxiety.
Parkinson's Disease (PD)	High-frequency rTMS may have beneficial effects on motor disorders
Alzheimer disease (AD)	High-frequency, offline rTMS may contribute to small short-term improvement in cognitive functioning

Table S2: Search syntax

AMED (Allied and Complementary Medicine) 1985 to May 2016		
#	Searches	Results
1	transcranial magnetic stimulation.mp.	287
2	TMS.mp.	116
3	Theory of mind.mp.	56
4	ToM.mp.	25
5	mentali*.mp.	20
6	role taking.mp.	3
7	perspective taking.mp.	5
8	empathy.mp.	343
9	1 or 2	313
10	3 or 4 or 5 or 6 or 7 or 8	429
11	9 and 10	1

Cochrane Library: Issue 4 of 12, April 2016; Cochrane Central Register of Controlled		
#	Searches	Results
#1	transcranial magnetic stimulation	2024
#2	TMS	796
#3	Theory of mind	659
#4	ToM	164
#5	mentali*	96
#6	role taking	800
#7	perspective taking	176
#8	empath*	453
#9	#1 or #2	2235
#10	#3 or #4 or #5 or #6 or #7 or #8	2233
#11	#9 and #10	6

OVID: Embase 1980 to 2016 Week 21		
#	Searches	Results
1	transcranial magnetic stimulation.mp.	18219
2	TMS.mp.	12740
3	Theory of mind.mp.	4908
4	ToM.mp.	3625
5	mentali*.mp.	3749
6	role taking.mp.	164
7	perspective taking.mp.	1354
8	empath*.mp.	23301
9	1 or 2	23283
10	3 or 4 or 5 or 6 or 7 or 8	33707
11	9 and 10	128

OVID MEDLINE(R) 1946 to May Week 2 2016		
#	Searches	Results
1	transcranial magnetic stimulation.mp.	10734
2	TMS.mp.	7672
3	Theory of mind.mp.	3010
4	ToM.mp.	2291
5	mentali*.mp.	2406
6	role taking.mp.	151
7	perspective taking.mp.	857
8	empath*.mp.	18755
9	1 or 2	13734
10	3 or 4 or 5 or 6 or 7 or 8	25376
11	9 and 10	59

OVID: PsycINFO 1806 to May Week 3 2016		
#	Searches	Results
1	transcranial magnetic stimulation.mp.	7371
2	TMS.mp.	3724
3	Theory of mind.mp.	7047
4	ToM.mp.	3343
5	mentali*.mp.	5698
6	role taking.mp.	2669
7	perspective taking.mp.	3265
8	empath*.mp.	26113
9	1 or 2	7824
10	3 or 4 or 5 or 6 or 7 or 8	42782
11	9 and 10	65

Pubmed 25052016		
#	Searches	Results
#1	Search (transcranial magnetic stimulation) OR TMS	16057
#2	Search (((((theory of mind) OR mentali*) OR empath*) OR perspective taking) OR role taking) OR ToM	61634
#3	Search (#1) AND #2	131

Web of Science Core Collection: Citation Indexes: Science Citation Index Expanded (SCI-EXPANDED) --1900-present; Social Sciences Citation Index (SSCI) --1956-present; Arts & Humanities Citation Index (A&HCI) --1975-present; Conference Proceedings Citation Index- Science (CPCI-S) --1990-present; Conference Proceedings Citation Index- Social Science & Humanities (CPCI-SSH) --1990-

present		
#	Searches	Results
#1	"transcranial magnetic stimulation"	16137
#2	TMS	13326
#3	"Theory of mind"	5489
#4	ToM	10802
#5	mentali*	6906
#6	"role taking"	436
#7	"perspective taking"	3171
#8	empath*	18938
#9	#1 or #2	23415
#10	#3 or #4 or #5 or #6 or #7 or #8	41869
#11	#9 and #10	116

Table S3: The list of the excluded studies

Excluded due to the type of publication

-
- Agnew, Z. K., Bhakoo, K. K., & Puri, B. K. (2007). The human mirror system: A motor resonance theory of mind-reading. *Brain Research Reviews*, 54(2), 286-293. doi: 10.1016/j.brainresrev.2007.04.003
-
- Andrews, S. C., Enticott, P. G., Hoy, K. E., & Fitzgerald, P. B. (2013). Mirror systems and social cognition in schizophrenia. *Schizophrenia Bulletin*, 39, S218. doi: <http://dx.doi.org/10.1093/schbul/sbt011>
-
- Avenanti, A. (2010). Neurophysiological markers of empathy for pain. *European Journal of Neurology*, 17, 10. doi: <http://dx.doi.org/10.1111/j.1468-1331.2010.03230.x>
-
- Avenanti, A., Candidi, M., & Urgesi, C. (2013). Vicarious motor activation during action perception: beyond correlational evidence. *Frontiers in Human Neuroscience*, 7. doi: 10.3389/fnhum.2013.00185
-
- Avenanti, A., & Urgesi, C. (2011). Understanding 'what' others do: mirror mechanisms play a crucial role in action perception. *Social Cognitive and Affective Neuroscience*, 6(3), 257-259. doi: 10.1093/scan/nsr004
-
- Baeken, C. (2011). One left-sided dorsolateral prefrontal cortical HF-rTMS session affects emotional neuronal processing in healthy women. *Clinical Neurophysiology*, 122, S144-S145. doi: <http://dx.doi.org/10.1016/S1388-2457%2811%2960516-6>
-
- Baeken, C., Van Schuerbeek, P., De Raedt, R., De Mey, J., Vanderhasselt, M. A., Santermans, L., . . . Luybaert, R. (2011). The effect of one left-sided prefrontal HF-rTMS session on emotional brain processes. *European Psychiatry*, 26. doi: <http://dx.doi.org/10.1016/S0924-9338%2811%2972838-3>
-
- Balconi, M., & Canavesio, Y. (2013). High-frequency rTMS stimulation improves the facial mimicry and detection responses in an empathic emotional task. *Clinical Neurophysiology*, 124 (10), e115-e116. doi: <http://dx.doi.org/10.1016/j.clinph.2013.04.184>
-
- Balconi, M., & Canavesio, Y. (2013). rTMS stimulation improves the facial mimicry and detection responses in an empathic emotional task. *Behavioural Neurology*, 27 (3), 418. doi: <http://dx.doi.org/10.3233/BEN-139900>
-
- Bernhardt, B. C., & Singer, T. (2012). The neural basis of empathy (pp. 1-23). 4139 El Camino Way, P.O. Box 10139, Palo Alto CA 94306, United States: Annual Reviews Inc.
-
- Bernier, R., & Dawson, G. (2009). The role of mirror neuron dysfunction in autism. *Mirror neuron systems: The Role of Mirroring Processes in Social Cognition* (pp. 261-286). Totowa, NJ: Humana Press; US.
-
- Bouaziz, N., Benadhira, R., Sidhoumi, D., & Januel, D. (2011). Transcranial magnetic stimulation (rTMS) concerning the treatment of schizophrenia: Interests and perspectives. *Annales Medico-Psychologiques*, 169(3), 192-195. doi: <http://dx.doi.org/10.1016/j.amp.2011.02.013>
-
- Christov-Moore, L., Simpson, E. A., Coude, G., Grigaityte, K., Iacoboni, M., & Ferrari, P. F. (2014). Empathy: Gender effects in brain and behavior. *Neuroscience and Biobehavioral Reviews*, 46(P4), 604-627. doi: <http://dx.doi.org/10.1016/j.neubiorev.2014.09.001>
-

-
- Cooper, N. R., Puzzo, I., & Pawley, A. D. (2008). Contagious yawning: The mirror neuron system may be a candidate physiological mechanism. *Medical Hypotheses*, 71(6), 975-976. doi: <http://dx.doi.org/10.1016/j.mehy.2008.07.023>
-
- Corbetta, M., Patel, G., & Shulman, G. L. (2008). The Reorienting System of the Human Brain: From Environment to Theory of Mind. *Neuron*, 58(3), 306-324. doi: <http://dx.doi.org/10.1016/j.neuron.2008.04.017>
-
- Demirtas-Tatlidede, A., & Schmahmann, J. D. (2013). Morality: Incomplete without the cerebellum? *Brain*, 136(8), e244. doi: <http://dx.doi.org/10.1093/brain/awt070>
-
- Enticott, P. G., Kennedy, H. A., Rinehart, N. J., May, S., Rossell, S., Tonge, B. J., . . . Fitzgerald, P. B. (2011). Social cognitive impairments in autism spectrum disorders: Insights from neuropsychiatry. *Clinical EEG and Neuroscience*, 42 (2), 130.
-
- Fumagalli, M., & Priori, A. (2012). Functional and clinical neuroanatomy of morality. *Brain*, 135(Pt 7), 2006-2021. doi: <http://dx.doi.org/10.1093/brain/awr334>
-
- Hetu, S., Taschereau-Dumouchel, V., & Jackson, P. L. (2012). Stimulating the brain to study social interactions and empathy. *Brain Stimulation*, 5(2), 95-102. doi: <http://dx.doi.org/10.1016/j.brs.2012.03.005>
-
- Iacoboni, M. (2012). The human mirror neuron system and its role in imitation and empathy *The primate mind: Built to connect with other minds* (pp. 32-47). Cambridge, MA: Harvard University Press; US.
-
- Iacoboni, M., & Dapretto, M. (2006). The mirror neuron system and the consequences of its dysfunction. *Nature Reviews Neuroscience*, 7(12), 942-951. doi: <http://dx.doi.org/10.1038/nrn2024>
-
- Jankowiak-Siuda, K., Siemieniuk, K., & Grabowska, A. (2009). Neurobiological basis of empathy. [Polish]
-
- Neurobiologiczne podstawy empatii. *Neuropsychiatria i Neuropsychologia*, 4(2), 51-58.
-
- Krippel, M., & Karim, A. A. (2011). "EuroTheory of mind" and its neuronal correlates in forensically relevant disorders. *Nervenarzt*, 82(7), 843-852. doi: 10.1007/s00115-010-3073-x
-
- Li, H., Wang, J., Li, C., & Xiao, Z. (2014). Repetitive transcranial magnetic stimulation (rTMS) for panic disorder in adults. *Cochrane Database of Systematic Reviews*, (9). <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD009083.pub2/abstract> doi:10.1002/14651858.CD009083.pub2
-
- Mak, A. D. P., & Lam, L. C. W. (2013). Neurocognitive profiles of people with borderline personality disorder. *Current Opinion in Psychiatry*, 26(1), 90-96. doi: <http://dx.doi.org/10.1097/YCO.0b013e32835b57a9>
-
- Mehta, U. M., Basavaraju, R., Thirthalli, J., & Gangadhar, B. N. (2012). Mirror neuron dysfunction-a neuro-marker for social cognition deficits in drug naive schizophrenia. *Biological Psychiatry*, 1), 314S. doi: <http://dx.doi.org/10.1016/j.biopsych.2012.02.014>
-
- Mehta, U. M., Basavaraju, R., Thirthalli, J., & Gangadhar, B. N. (2013). Mirror neuron dysfunction in schizophrenia and its association with social cognition. *Schizophrenia Bulletin*, 39, S242. doi: <http://dx.doi.org/10.1093/schbul/sbt011>
-
- Miniussi, C., Cappa, S. F., Cohen, L. G., Floel, A., Fregni, F., Nitsche, M. A., . . . Walsh, V. (2008). Efficacy of repetitive transcranial magnetic
-

stimulation/transcranial direct current stimulation in cognitive neurorehabilitation. *Brain Stimul*, 1(4), 326-336. doi: 10.1016/j.brs.2008.07.002

Molnar-Szakacs, I. (2011). From actions to empathy and morality - A neural perspective. *Journal of Economic Behavior & Organization*, 77(1), 76-85. doi: 10.1016/j.jebo.2010.02.019

Newlin, D. B., & Renton, R. M. (2010). A Self in the Mirror: Mirror Neurons, Self-Referential Processing, and Substance Use Disorders. *Substance Use and Misuse*, 45(11), 1697-1726. doi: 10.3109/10826084.2010.482421

Obhi, S. S., & Sebanz, N. (2011). Moving together: Toward understanding the mechanisms of joint action. *Experimental Brain Research*, 211(3-4), 329-336. doi: <http://dx.doi.org/10.1007/s00221-011-2721-0>

O'Malley, M. K., Ro, T., & Levin, H. S. (2006). Assessing and inducing neuroplasticity with transcranial magnetic stimulation and robotics for motor function. *Archives of Physical Medicine and Rehabilitation*, 87(12 Suppl 2), S59-66. doi: 10.1016/j.apmr.2006.08.332

Perkins, T., Stokes, M., McGillivray, J., & Bittar, R. (2010). Mirror neuron dysfunction in autism spectrum disorders. *Journal of Clinical Neuroscience*, 17(10), 1239-1243. doi: <http://dx.doi.org/10.1016/j.jocn.2010.01.026>

Schuerker, T., Langguth, B., & Sommer, M. (2014). Modulating functional and dysfunctional mentalizing by transcranial magnetic stimulation. *Frontiers in Psychology*, 5. doi: 10.3389/fpsyg.2014.01309

Singer, T., & Frith, C. (2005). The painful side of empathy. *Nature Neuroscience*, 8(7), 845-846. doi: <http://dx.doi.org/10.1038/nn0705-845>

Suttrup, J., Keysers, C., & Thioux, M. (2015). The role of the theory of mind network in action observation-an rTMS study. *Brain Stimulation*, 8 (2), 415-416.

van Honk, J., & Schutter, D. J. (2006). Unmasking feigned sanity: A neurobiological model of emotion processing in primary psychopathy. *Cognitive Neuropsychiatry*, 11(3), 285-306. doi: <http://dx.doi.org/10.1080/13546800500233728>

Excluded due to no TMS involved after reviewing abstracts

Aziz-Zadeh, L., Sheng, T., & Gheyntanchi, A. (2010). Common Premotor Regions for the Perception and Production of Prosody and Correlations with Empathy and Prosodic Ability. *PloS One*, 5(1). doi: 10.1371/journal.pone.0008759

Benuzzi, F., Lui, F., Duzzi, D., Nichelli, P. F., & Porro, C. A. (2008). Does it look painful or disgusting? Ask your parietal and cingulate cortex. *Journal of Neuroscience*, 28(4), 923-931. doi: 10.1523/jneurosci.4012-07.2008

Lepage, J.-F. (2011). *Developpement et fonctionnement des mecanismes de resonance motrice chez l'humain*. Dissertation Abstracts International: Section B: The Sciences and Engineering, 72(4-B), 2475.

Marsh, L. E., Mullett, T. L., Ropar, D., & de, C. (2014). Responses to irrational actions in action observation and mentalising networks of the human brain. *Neuroimage*, 103, 81-90. doi: <http://dx.doi.org/10.1016/j.neuroimage.2014.09.020>

Parkinson, C., & Wheatley, T. (2014). Relating Anatomical and Social Connectivity: White Matter Microstructure Predicts Emotional Empathy. *Cerebral Cortex*, 24(3), 614-625. doi: 10.1093/cercor/bhs347

Excluded due to intervention (not rTMS) after reviewing abstracts

-
- Andrews, S. C., Enticott, P. G., Hoy, K. E., Thomson, R. H., & Fitzgerald, P. B. (2015). No evidence for mirror system dysfunction in schizophrenia from a multimodal TMS/EEG study. *Psychiatry Research*. doi: 10.1016/j.psychres.2015.05.067
-
- Andrews, S. C., Enticott, P. G., Hoy, K. E., Thomson, R. H., & Fitzgerald, P. B. (2015). Reduced mu suppression and altered motor resonance in euthymic bipolar disorder: Evidence for a dysfunctional mirror system? *Social Neuroscience*, 1-12. doi: 10.1080/17470919.2015.1029140
-
- Bolognini, N., Rossetti, A., Fusaro, M., Vallar, G., & Miniussi, C. (2014). Sharing social touch in the primary somatosensory cortex. *Current Biology*, 24(13), 1513-1517. doi: 10.1016/j.cub.2014.05.025
-
- Borgomaneri, S., Gazzola, V., & Avenanti, A. (2012). Motor mapping of implied actions during perception of emotional body language. *Brain Stimulation*, 5(2), 70-76. doi: 10.1016/j.brs.2012.03.011
-
- Borgomaneri, S., Gazzola, V., & Avenanti, A. (2014). Temporal dynamics of motor cortex excitability during perception of natural emotional scenes. *Social Cognitive and Affective Neuroscience*, 9(10), 1451-1457. doi: 10.1093/scan/nst139
-
- Fourkas, A. D., Avenanti, A., Urgesi, C., & Aglioti, S. M. (2006). Corticospinal facilitation during first and third person imagery. *Experimental Brain Research*, 168(1-2), 143-151. doi: <http://dx.doi.org/10.1007/s00221-005-0076-0>
-
- Lepage, J. F., Tremblay, S., & Theoret, H. (2010). Early non-specific modulation of corticospinal excitability during action observation. *European Journal of Neuroscience*, 31(5), 931-937. doi: <http://dx.doi.org/10.1111/j.1460-9568.2010.07121.x>
-
- Liuzza, M. T., Candidi, M., Sforza, A. L., & Aglioti, S. M. (2015). Harm avoiders suppress motor resonance to observed immoral actions. *Social Cognitive and Affective Neuroscience*, 10(1), 72-77. doi: 10.1093/scan/nsu025
-
- Mahayana, I. T., Banissy, M. J., Chen, C.-Y., Walsh, V., Juan, C.-H., & Muggleton, N. G. (2014). Motor empathy is a consequence of misattribution of sensory information in observers. *Frontiers in Human Neuroscience*, 8, 47.
-
- Minio-Paluello, I., Baron-Cohen, S., Avenanti, A., Walsh, V., & Aglioti, S. M. (2009). Absence of Embodied Empathy During Pain Observation in Asperger Syndrome. *Biological Psychiatry*, 65(1), 55-62. doi: <http://dx.doi.org/10.1016/j.biopsych.2008.08.006>
-

Excluded due to outcome (not measuring empathy or ToM) after reviewing abstracts

-
- Basavaraju, R., Mehta, U. M., & Thirhalli, J. (2014). Mirror neuron activity and symptom dimensions in drug-naive mania-a transcranial magnetic stimulation study. *Bipolar Disorders*, 16, 83-84. doi: <http://dx.doi.org/10.1111/bdi.12189>
-
- Bolognini, N., Rossetti, A., Maravita, A., & Miniussi, C. (2011). Seeing Touch in the Somatosensory Cortex: ATMS Study of the Visual Perception of Touch. *Human Brain Mapping*, 32(12), 2104-2114. doi: 10.1002/hbm.21172
-
- Brune, M., Scheele, D., Heinisch, C., Tas, C., Wischniewski, J., & Gunturkun, O. (2012). Empathy Moderates the Effect of Repetitive Transcranial Magnetic Stimulation of the Right Dorsolateral Prefrontal Cortex on Costly Punishment. *PloS One*, 7(9). doi: <http://dx.doi.org/10.1371/journal.pone.0044747>
-

Catmur, C., Walsh, V., & Heyes, C. (2007). Sensorimotor learning configures the human mirror system. *Current Biology*, 17(17), 1527-1531. doi: 10.1016/j.cub.2007.08.006

Cazzato, V., Mian, E., Serino, A., Mele, S., & Urgesi, C. (2015). Distinct contributions of extrastriate body area and temporoparietal junction in perceiving one's own and others' body. *Cognitive Affective & Behavioral Neuroscience*, 15(1), 211-228. doi: 10.3758/s13415-014-0312-9

Chiang, T. C., Lu, R. B., Hsieh, S., Chang, Y. H., & Yang, Y. K. (2014). Stimulation in the dorsolateral prefrontal cortex changes subjective evaluation of percepts. *PloS One*, 9(9), e106943. doi: 10.1371/journal.pone.0106943

Du, D. Q., & Wu, Y. B. (2005). Living ability and cognitive function ameliorated by low frequency repetitive transcranial magnetic stimulation in patients with post-stroke depression: Comparison with drug plus psychological treatment. [Chinese]. *Chinese Journal of Clinical Rehabilitation*, (16), 22-23. <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/568/CN-00557568/frame.html>

Fitzgibbon, B. M., Enticott, P. G., Bradshaw, J. L., Giummarra, M. J., Chou, M., Georgiou-Karistianis, N., & Fitzgerald, P. B. (2012). Enhanced corticospinal response to observed pain in pain synesthetes. *Cognitive, Affective and Behavioral Neuroscience*, 12(2), 406-418. doi: <http://dx.doi.org/10.3758/s13415-011-0080-8>

Knoch, D., Gianotti, L. R., Pascual-Leone, A., Treyer, V., Regard, M., Hohmann, M., & Brugger, P. (2006). Disruption of right prefrontal cortex by low-frequency repetitive transcranial magnetic stimulation induces risk-taking behavior. *Journal of Neuroscience*, 26(24), 6469-6472. doi: 10.1523/jneurosci.0804-06.2006

Novembre, G., Ticini, L., Schutz-Bosbach, S., & Keller, P. (2013). Motor simulation coordinates joint actions in real time: Music performance meets on-line double-pulse TMS. *Clinical Neurophysiology*, 124 (10), e82. doi: <http://dx.doi.org/10.1016/j.clinph.2013.04.116>

Novembre, G., Ticini, L. F., Schutz-Bosbach, S., & Keller, P. E. (2012). Distinguishing self and other in joint action. Evidence from a musical paradigm. *Cerebral Cortex*, 22(12), 2894-2903. doi: <http://dx.doi.org/10.1093/cercor/bhr364>

Papeo, L., Corradi-Dell'Acqua, C., & Rumiati, R. I. (2011). "She" Is Not Like "I": The Tie between Language and Action Is in Our Imagination. *Journal of Cognitive Neuroscience*, 23(12), 3939-3948.

Pretalli, J. B., Nicolier, M., Chopard, G., Vandell, P., Tio, G., Monnin, J., . . . Haffen, E. (2012). Resting motor threshold changes and clinical response to prefrontal repetitive transcranial magnetic stimulation in depressed patients. *Psychiatry and Clinical Neurosciences*, 66(4), 344-352. doi: 10.1111/j.1440-1819.2012.02341.x

White, N. C., Reid, C., & Welsh, T. N. (2014). Responses of the human motor system to observing actions across species: A transcranial magnetic stimulation study. *Brain and Cognition*, 92, 11-18. doi: <http://dx.doi.org/10.1016/j.bandc.2014.10.004>

Excluded due to intervention (not rTMS) after reviewing full-texts

Avenanti, A., & Aglioti, S. M. (2006). Pain in the motor system: One study of transcranial magnetic stimulation. *Giornale Italiano di Psicologia*, 33(4), 777-792.

-
- Avenanti, A., Buetti, D., Galati, G., & Aglioti, S. M. (2005). Transcranial magnetic stimulation highlights the sensorimotor side of empathy for pain. *Nature Neuroscience*, 8(7), 955-960. doi: 10.1038/nn1481
-
- Avenanti, A., Minio-Paluello, I., Bufalari, I., & Aglioti, S. M. (2006). Stimulus-driven modulation of motor-evoked potentials during observation of others' pain. *Neuroimage*, 32(1), 316-324. doi: 10.1016/j.neuroimage.2006.03.010
-
- Avenanti, A., Minio-Paluello, I., Bufalari, I., & Aglioti, S. M. (2009). The pain of a model in the personality of an onlooker: Influence of state-reactivity and personality traits on embodied empathy for pain. *Neuroimage*, 44(1), 275-283. doi: <http://dx.doi.org/10.1016/j.neuroimage.2008.08.001>
-
- Avenanti, A., Minio-Paluello, I., Sforza, A., & Aglioti, S. M. (2009). Freezing or escaping? Opposite modulations of empathic reactivity to the pain of others. *Cortex*, 45(9), 1072-1077. doi: <http://dx.doi.org/10.1016/j.cortex.2008.10.004>
-
- Avenanti, A., Sirigu, A., & Aglioti, S. M. (2010). Racial bias reduces empathic sensorimotor resonance with other-race pain. *Current Biology*, 20(11), 1018-1022. doi: <http://dx.doi.org/10.1016/j.cub.2010.03.071>
-
- Borgomaneri, S., Gazzola, V., & Avenanti, A. (2014). Transcranial magnetic stimulation reveals two functionally distinct stages of motor cortex involvement during perception of emotional body language. *Brain Structure & Function* Jul(Pagination), No Pagination Specified. doi: <http://dx.doi.org/10.1007/s00429-014-0825-6>
-
- De Coster, L., Andres, M., & Brass, M. (2014). Effects of being imitated on motor responses evoked by pain observation: Exerting control determines action tendencies when perceiving pain in others. *The Journal of Neuroscience*, 34(20), 6952-6957. doi: <http://dx.doi.org/10.1523/JNEUROSCI.5044-13.2014>
-
- Donne, C. M., Enticott, P. G., Rinehart, N. J., & Fitzgerald, P. B. (2011). A transcranial magnetic stimulation study of corticospinal excitability during the observation of meaningless, goal-directed, and social behaviour. *Neuroscience Letters*, 489(1), 57-61. doi: 10.1016/j.neulet.2010.11.067
-
- Enticott, P. G., Harrison, B. A., Arnold, S. L., Nibaldi, K., Segrave, R. A., Fitzgibbon, B. M., . . . Fitzgerald, P. B. (2012). Emotional valence modulates putative mirror neuron activity. *Neuroscience Letters*, 508(1), 56-59. doi: 10.1016/j.neulet.2011.12.018
-
- Enticott, P. G., Johnston, P. J., Herring, S. E., Hoy, K. E., & Fitzgerald, P. B. (2008). Mirror neuron activation is associated with facial emotion processing. *Neuropsychologia*, 46(11), 2851-2854. doi: 10.1016/j.neuropsychologia.2008.04.022
-
- Enticott, P. G., Kennedy, H. A., Bradshaw, J. L., Rinehart, N. J., & Fitzgerald, P. B. (2010). Understanding mirror neurons: Evidence for enhanced corticospinal excitability during the observation of transitive but not intransitive hand gestures. *Neuropsychologia*, 48(9), 2675-2680. doi: 10.1016/j.neuropsychologia.2010.05.014
-
- Enticott, P. G., Kennedy, H. A., Bradshaw, J. L., Rinehart, N. J., & Fitzgerald, P. B. (2011). Motor corticospinal excitability during the observation of interactive hand gestures. *Brain Research Bulletin*, 85(3-4), 89-95. doi: <http://dx.doi.org/10.1016/j.brainresbull.2011.03.018>
-
- Fecteau, S., Pascual-Leone, A., & Theoret, H. (2008). Psychopathy and the mirror neuron system: preliminary findings from a non-psychiatric sample.
-

Psychiatry Research, 160(2), 137-144. doi:

<http://dx.doi.org/10.1016/j.psychres.2007.08.022>

Fitzgibbon, B. M., Enticott, P. G., Bradshaw, J. L., Giummarra, M. J., Georgiou-Karistianis, N., Chou, M., & Fitzgerald, P. B. (2012). Motor cortical excitability and inhibition in acquired mirror pain. *Neuroscience Letters*, 530(2), 161-165. doi:

<http://dx.doi.org/10.1016/j.neulet.2012.09.036>

Guise, K., Kelly, K., Romanowski, J., Vogeley, K., Platek, S. M., Murray, E., & Keenan, J. P. (2007). The anatomical and evolutionary relationship between self-awareness and Theory of mind. *Human Nature*, 18(2), 132-142. doi:

<http://dx.doi.org/10.1007/s12110-007-9009-x>

Lepage, J. F., Lortie, M., Deal, C. L., & Theoret, H. (2014). Empathy, autistic traits, and motor resonance in adults with Turner syndrome. *Social Neuroscience*, 9(6), 601-609. doi: Doi 10.1080/17470919.2014.944317

Mehta, U. M., Thirthalli, J., Basavaraju, R., Gangadhar, B. N., & Pascual-Leone, A. (2014). Reduced mirror neuron activity in schizophrenia and its association with theory of mind deficits: Evidence from a transcranial magnetic stimulation study. *Schizophrenia Bulletin*, 40(5), 1083-1094. doi:

<http://dx.doi.org/10.1093/schbul/sbt155>

Minio-Paluello, I., Avenanti, A., & Aglioti, S. M. (2006). Left hemisphere dominance in reading the sensory qualities of others' pain? *Social Neuroscience*, 1(3-4), 320-333.

Wood, R., Gallese, V., & Cattaneo, L. (2010). Visuotactile empathy within the primary somatosensory cortex revealed by short-latency afferent inhibition. *Neuroscience Letters*, 473(1), 28-31. doi:

<http://dx.doi.org/10.1016/j.neulet.2010.02.012>

Excluded due to outcome (not measuring empathy or ToM) after reviewing full-texts

Baeken, C., Van Schuerbeek, P., De Raedt, R., De Mey, J., Vanderhasselt, M. A., Bossuyt, A., & Luybaert, R. (2011). The effect of one left-sided dorsolateral prefrontal sham-controlled HF-rTMS session on approach and withdrawal related emotional neuronal processes. *Clinical Neurophysiology*, 122(11), 2217-2226. doi: <http://dx.doi.org/10.1016/j.clinph.2011.04.009>

Brune, M., Scheele, D., Heinisch, C., Tas, C., Wischniewski, J., & Gunturkun, O. (2012). Empathy Moderates the Effect of Repetitive Transcranial Magnetic Stimulation of the Right Dorsolateral Prefrontal Cortex on Costly Punishment. *PloS One*, 7(9). doi: <http://dx.doi.org/10.1371/journal.pone.0044747>

Cazzato, V., Mian, E., Serino, A., Mele, S., & Urgesi, C. (2015). Distinct contributions of extrastriate body area and temporoparietal junction in perceiving one's own and others' body. *Cognitive Affective & Behavioral Neuroscience*, 15(1), 211-228. doi: 10.3758/s13415-014-0312-9

David, N., Jansen, M., Cohen, M. X., Osswald, K., Molnar-Szakacs, I., Newen, A., . . . Paus, T. (2009). Disturbances of self-other distinction after stimulation of the extrastriate body area in the human brain. *Social Neuroscience*, 4(1), 40-48. doi: 10.1080/17470910801938023

Rossetti, A., Miniussi, C., Maravita, A., & Bolognini, N. (2012). Visual perception of bodily interactions in the primary somatosensory cortex. *European Journal of Neuroscience*, 36(3), 2317-2323. doi: 10.1111/j.1460-9568.2012.08137.x

Rossetti, A., Miniussi, C., Maravita, A., & Bolognini, N. (2012). Visual perception of bodily interactions in the primary somatosensory cortex. *European Journal of Neuroscience*, 36(3), 2317-2323. doi: 10.1111/j.1460-9568.2012.08137.x

Rossetti, A., Miniussi, C., Maravita, A., & Bolognini, N. (2012). Visual perception of bodily interactions in the primary somatosensory cortex. *European Journal of Neuroscience*, 36(3), 2317-2323. doi: 10.1111/j.1460-9568.2012.08137.x

Table S4. Component and overall quality ratings of the reviewed studies

Study	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Overall
Balconi & Bortolotti, 2012	+	+++	+++	++	+++	+++	++
Balconi & Bortolotti, 2013	+	+++	+++	++	+++	+++	++
Balconi, Bortolotti, & Gonzaga, 2011	+	+++	+++	++	+++	+++	++
Balconi & Canavesio, 2013	+	+++	+++	++	+++	+++	++
Balconi & Canavesio, 2016	+	+++	+++	++	+++	+++	++
Balconi, Crivelli, & Bortolotti, 2010	+	+++	+++	++	+++	+++	++
Bolognini et al., 2013	+	+++	+++	++	+++	+++	++
Costa et al., 2008	+	+++	+++	++	+++	+++	++
Enticott et al., 2014	++	+++	+++	+++	+++	+++	+++
Giardina et al., 2011	+	+++	+++	++	+++	+++	++
Hoekert et al., 2010,	+	+++	+++	++	+++	+++	++
Kalbe et al., 2010	+	+++	+++	++	+++	+++	++
Keuken et al., 2011	+	+++	+++	++	+++	+++	++
Krause et al., 2012	+	+++	+++	++	+++	+++	++
Lev-Ran et al., 2012	+	+++	+++	++	+++	+++	++
Michael et al., 2014	+	+++	+++	++	+	+++	+
Pobric and Hamilton, 2006	+	+++	+++	++	+++	+++	++

Schuwerk et al., 2014	+	+++	+++	++	+++	+++	++
Silani et al., 2013	+	+++	+	++	+++	+++	+
Uddin et al., 2006	+	+++	+++	++	++	+++	++
Young et al., 2010	+	+++	+++	++	+++	+++	++

+ = weak, ++ = moderate, +++ = strong




Figure S1. Study Selection and Search Results




Figure S2a. Funnel plot of cognitive ToM trials included in the meta-analysis




Figure S2b. Filled funnel plot of the cognitive ToM trials in the meta-analysis after trim procedure

Empty dots with an outer square represent imputed missing trials.




Figure S3a. Funnel plot of the affective ToM trials in the meta-analysis




Figure S3b. Filled funnel plot of the affective ToM trials in the meta-analysis after trim procedure

No missing trials were found.